


MATH 5143 - Lecture # 22



MATH 5143 - Lecture 22

Clarifications about universal enveloping algebras

L Lie algebra

$T(L)$ tensor algebra (assoc.) alg. but noncommutative

$S(L)$ symmetric algebra (assoc.) alg. that is commutative

$U(L)$ univ. enveloping algebra (assoc.) alg. also NOT commutative

What's true: the associated graded algebra of $U(L)$ is $\cong S(L)$

in $U(L)$ we have $\underbrace{XY}_{\text{deg } 2} = \underbrace{YX}_{\text{deg } 2} + \underbrace{[X, Y]}_{\text{deg } 1} \neq YX$

key point: multiplication of homogeneous tensors in $U(L)$
IGNORING LOWER ORDER TERMS is commutative

Representation theory setup: L is a semisimple fin. dim. Lie algebra / \mathbb{F}

$\mathfrak{H} \subseteq L$ is a Cartan subalgebra, $\Phi \subset \mathfrak{H}^*$ is the root system,

$\Delta = \{\alpha_1, \alpha_2, \dots, \alpha_n\} \subset \Phi$ is a chosen base, $\Phi^+ = \{\text{positive roots}\}$,

and $W = \langle r_\alpha \mid \alpha \in \Phi \rangle = \langle r_\alpha \mid \alpha \in \Delta \rangle \subseteq GL(\mathfrak{H}^*)$

First key observation: Any finite-dim L -module V decomposes as a

direct sum of weight spaces $V = \bigoplus_{\lambda \in \mathfrak{H}^*} V_\lambda$ where $V_\lambda \stackrel{\text{def}}{=} \{v \in V \mid h \cdot v = \lambda(h)v \ \forall h \in \mathfrak{H}\}$

Call $\lambda \in \mathfrak{H}^*$ a weight of V if $V_\lambda \neq 0$ and call V_λ a weight space.

We make some definitions if $\dim V = \infty$ but in that case

the sum of weight spaces $\bigoplus_{\lambda \in \mathfrak{H}^*} V_\lambda$ [which is always direct] may be $\subsetneq V$.

Idea: to avoid pathologies with arbitrary infinite-dim L -modules, we consider standard cyclic modules.

A maximal vector in an arbitrary L -module V is a nonzero element v^+ in some weight space of V with $xv^+ = 0 \quad \forall x \in L_\alpha \quad \forall \alpha \in \Delta$

Lie's theorem (applied to the Borel algebra $\mathfrak{B} = \mathfrak{H} \oplus \bigoplus_{\alpha \in \Phi^+} L_\alpha$ acting on V) ensures that V has a maximal vector whenever $\dim V < \infty$.

Def A standard cyclic module for L is an L -module V

with a maximal vector v^+ such that $V = \mathcal{U}(L) \cdot v^+$.

In this situation, the maximal vector v^+ will belong to V_λ

for some $\lambda \in \mathfrak{H}^*$, and we say V has highest weight λ
and we call v^+ is a highest weight vector

Thm (Structure thm for standard cyclic modules)

For each $\beta \in \Phi^+$ choose $x_\beta \in L_\beta, y_\beta \in L_{-\beta}$ such that $[x_\beta, y_\beta] = h_\beta \in H$

Write $\lambda > \mu$ for $\lambda, \mu \in H^*$ if $\lambda - \mu \in \mathbb{Z}_{\geq 0} \cdot \text{span}\{\alpha \in \Delta\}$

Suppose V is standard cyclic L -module with maximal vector $v^+ \in V_\lambda$

(a) If $\Phi^+ = \{\beta_1, \beta_2, \beta_3, \dots\}$ then V is spanned by vectors of the form

$$y_{\beta_{i_1}} y_{\beta_{i_2}} y_{\beta_{i_3}} \dots y_{\beta_{i_k}} v^+ \quad \text{where } 1 \leq i_1 \leq i_2 \leq i_3 \leq \dots \leq i_k$$

(b) All weights μ for V have $\mu < \lambda$ and $\dim V_\mu < \infty$ and $\dim V_\lambda = 1$.

(c) Any submodule of V is a direct sum of its weight spaces

(d) V is indecomposable with unique maximal proper submodule and unique irreducible quotient

(e) Every homomorphic image of V is standard cyclic of same weight λ

Today: first, two more thms about standard cyclic modules

Thm A If V and W are irreducible standard cyclic L -modules with same highest weight $\lambda \in \mathfrak{H}^*$ then $V \cong W$

Thm B If $\lambda \in \mathfrak{H}^*$ then there exists an irreducible standard cyclic L -module $V(\lambda)$ of highest weight λ .

Pf of Thm A. Let $X = V \oplus W \stackrel{\text{def}}{=} \{v+w \mid v \in V, w \in W\}$. This is an L -module and if $v^+ \in V$ and $w^+ \in W$ are highest weight vectors then $x^+ \stackrel{\text{def}}{=} v^+ + w^+ \in X$ is a maximal vector also of weight λ .

Let Y be the submodule of X generated by x^+ . This is standard cyclic by def.

But $V \cong Y / \ker \pi_1$ and $W \cong Y / \ker \pi_2$ where $\pi_1: Y \rightarrow V$ and $\pi_2: Y \rightarrow W$ are the obvious surjective homomorphisms. This means V and W are both isomorphic to the unique irreducible quotient of Y . \square

To prove Thm B we must explain how to construct standard $U(\mathfrak{g})$ modules

Induced modules Begin with a 1-dim vector space $D_\lambda = \mathbb{F}\text{span}\{v^+\}$ spanned by some vector v^+ . Let $\lambda \in \mathfrak{H}^*$ and $\mathfrak{B} = \mathfrak{B}(\lambda) \stackrel{\text{def}}{=} \mathfrak{H} \oplus \bigoplus_{\alpha \in \Phi^+} \mathfrak{L}_\alpha \subset \mathfrak{L}$.

The Borel subalgebra \mathfrak{B} acts on D_λ linearly by

$$h \cdot v^+ \stackrel{\text{def}}{=} \lambda(h)v^+ \quad \text{and} \quad x v^+ \stackrel{\text{def}}{=} 0 \quad \text{for } h \in \mathfrak{H}, \alpha \in \Phi^+, x \in \mathfrak{L}_\alpha$$

This makes D_λ into a module for \mathfrak{B} and for $U(\mathfrak{B})$

Def Let $Z(\lambda) \stackrel{\text{def}}{=} U(\mathfrak{L}) \otimes_{U(\mathfrak{B})} D_\lambda$

This is a general construction of a $U(\mathfrak{L})$ -module: $U(\mathfrak{L})$ (left $U(\mathfrak{L})$ -mod) $+$ D_λ (left $U(\mathfrak{B})$ -mod)

Concretely, $Z(\lambda)$ is vector space spanned by

the tensors $x \otimes y$ ($x \in U(\mathfrak{L}), y \in D_\lambda$) subject to relations

$$(x+x') \otimes y = x \otimes y + x' \otimes y \quad c(x \otimes y) = (cx) \otimes y = x \otimes (cy)$$

$$x \otimes (y+y') = x \otimes y + x \otimes y' \quad xb \otimes y = x \otimes by \quad \text{for } x \in U(\mathfrak{L}), b \in U(\mathfrak{B}), y \in D_\lambda$$

The way L acts on $Z(\lambda)$ is $A \cdot (x \otimes y) \stackrel{\text{def}}{=} (Ax) \otimes y$ ($w /$ highest weight vector $\nearrow 1 \otimes v^+$)

Claim $Z(\lambda)$ is a standard cyclic L -module of weight λ .

Pf Every $y \in D_\lambda$ is a scalar multiple of v^+ so every tensor

$x \otimes y \in Z(\lambda)$ is equal to $\tilde{x} \cdot (1 \otimes v^+)$ where $\tilde{x} \in U(L)$ is a scalar

multiple of x . For $x \in L_\alpha$ for $\alpha \in \Delta$ we have $x \in \mathfrak{B}$ so

$$x \cdot (1 \otimes v^+) = x \otimes v^+ = 1 \otimes xv^+ = 1 \otimes 0 = 0. \text{ Also for } h \in \mathfrak{H} \subset \mathfrak{B}$$

$$\text{we have } h \cdot (1 \otimes v^+) = h \otimes v^+ = 1 \otimes hv^+ = 1 \otimes \lambda(h)v^+ = \lambda(h)(1 \otimes v^+) \square$$

Let $N^- = \bigoplus_{\alpha \in -\Phi^+} L_\alpha$. The relation $x b \otimes v^+ = x \otimes b v^+ \quad \forall b \in \mathfrak{B}$, since $L = \mathfrak{N}^- \oplus \mathfrak{B}$,

implies that if $\bar{\Phi}^+ = \{\beta_1, \beta_2, \beta_3, \dots\}$ and $\left. \begin{array}{l} x_i = x_{\beta_i} \text{ spans } L_{\beta_i} \\ y_i = y_{\beta_i} \text{ spans } L_{-\beta_i} \end{array} \right\}$ then

$$\left\{ y_{i_1} y_{i_2} \dots y_{i_k} \otimes v^+ \mid k \geq 0 \text{ and } i_1 \leq i_2 \leq \dots \leq i_k \right\}$$

is a basis for $Z(\lambda)$, via the PBW theorem.

Prop $Z(\lambda) \cong U(L) / I(\lambda)$ as $U(L)$ -modules, where $I(\lambda)$ is left ideal generated in $U(L)$ by the elements

$$\{x_1, x_2, x_3, \dots\} \cup \{h_\alpha - \lambda(h_\alpha) \cdot 1 \mid \alpha \in \bar{\Phi}^+\}$$

Pf These generators annihilate $1 \otimes v^+$ so there is a surjective morphism $U(L) / I(\lambda) \rightarrow Z(\lambda)$ which is injective using PBW theorem. \square

Thm (Thm B) Define $V(\lambda)$ for $\lambda \in H^*$ to be the unique irreducible quotient of the standard cyclic module $Z(\lambda)$.

Then $V(\lambda)$ is standard cyclic of weight λ and irreducible.

Note: $V(\lambda)$ still might be infinite-dimensional

Pf Since $Z(\lambda)$ is standard cyclic, and since $V(\lambda)$ as a quotient is a homomorphic image of $Z(\lambda)$, every thing follows from structure theorem for standard cyclic modules. \square

In some sense, hardest part of thm is showing $Z(\lambda) \neq 0$
(but we will not discuss this issue in detail, follows from PBW thm)

Two new goals: ① Explain when $V(\lambda)$ is finite dim.

② Determine weight spaces $V(\lambda)_\mu \subseteq V(\lambda)$

Fact If V is any irreducible L -module with $\dim V < \infty$
then $V \cong V(\lambda)$ for some $\lambda \in \mathfrak{h}^*$.

Pf If $\dim V < \infty$ then Lie's thm applied to B -action on V
implies existence of a maximal vector of some weight λ .

This vector must generate V by irreducibility, so $V \cong V(\lambda)$ by Thm A. \square

To be continued next time.