

This document is a **transcript** of the lecture, with extra summary and vocabulary sections for your convenience. Due to time constraints, the notes may sometimes only contain limited illustrations, proofs, and examples; for a more thorough discussion of the course content, **consult the textbook**.

Summary

Quick summary of today's notes. Lecture starts on next page.

- The determinant has a geometric interpretation in terms of volume. This is the reason why determinants appear when you do substitutions in multivariable integrals.

The columns of a 2×2 matrix A are the sides of a unique parallelogram in \mathbb{R}^2 .

The absolute value of $\det A$ is the area of this parallelogram.

This fact generalizes to n dimensions if we replace “parallelogram” by its n -dimensional analogue.

- We introduce the concept of a *vector space* to generalize the idea of a subspace of \mathbb{R}^n .

Formally, an (*abstract*) *vector space* is a nonempty set with a “zero vector” and two operations that can be thought of a “vector addition” and “scalar multiplication.”

These operations are subject to several conditions.

All subspaces of \mathbb{R}^n , including \mathbb{R}^n itself, are examples of vector spaces.

The set of polynomials in one variable is another example of a vector space.

- There are notions of linear combinations, span, linear independence, subspaces, bases, and dimension for vector spaces. The definitions are the same as the ones we already used for \mathbb{R}^n .
- If X and Y are sets, then let $\text{Functions}(X, Y)$ be the set of functions $f : X \rightarrow Y$.

The sets $\text{Functions}(X, \mathbb{R})$ and $\text{Functions}(X, \mathbb{R}^n)$ are vector spaces.

More generally, if V is a vector space, then $\text{Functions}(X, V)$ is a vector space.

The corresponding vector operations and zero vector are

$$\begin{aligned} f + g &= (\text{ the function that maps } x \mapsto f(x) + g(x) \text{ for } x \in X), \\ cf &= (\text{ the function that maps } x \mapsto c \cdot f(x) \text{ for } x \in X), \\ 0 &= (\text{ the function that maps } x \mapsto 0 \in V \text{ for } x \in X), \end{aligned}$$

for $f, g \in \text{Functions}(X, V)$ and $c \in \mathbb{R}$.

Most abstract vector spaces of interest arise as subspaces of $\text{Functions}(X, V)$ for some V .

- If U and V are vector spaces then a function $f : U \rightarrow V$ is *linear* if

$$f(u + v) = f(u) + f(v) \quad \text{and} \quad f(cv) = c \cdot f(v)$$

for all $u, v \in U$ and $c \in \mathbb{R}$.

1 Last time: determinants

Let n be a positive integer.

Theorem. The determinant is the unique function $\det : \{ n \times n \text{ matrices} \} \rightarrow \mathbb{R}$ such that

$$(1) \det I_n = 1 \text{ where } I_n = \begin{bmatrix} 1 & & & \\ & \ddots & & \\ & & \ddots & \\ & & & 1 \end{bmatrix} \text{ is the } n \times n \text{ identity matrix.}$$

(2) Switching two columns reverses the sign of the determinant.

(3) $\det A$ is linear as a function of a single column A if all other columns are fixed.

For 1×1 and 2×2 matrices, we have $\det [a] = a$ and $\det \begin{bmatrix} a & b \\ c & d \end{bmatrix} = ad - bc$.

The *diagonal (positions)* of an $n \times n$ matrix are the positions $(1, 1), (2, 2), \dots, (n, n)$.

The *diagonal entries* of a matrix are the entries in these positions.

A matrix is *upper triangular* if all of its nonzero entries are in positions on or above the diagonal.

A matrix is *lower triangular* if all of its nonzero entries are in positions on or below the diagonal.

A *triangular matrix* is a square matrix that is either upper or lower triangular.

A *diagonal matrix* is a matrix that is both upper and lower triangular: in other words, all of its nonzero entries appear in diagonal positions.

Theorem. If A is triangular square matrix then $\det A$ is the product of the diagonal entries of A .

Theorem. A square matrix A is invertible if and only if $\det A \neq 0$.

Theorem. If A and B are $n \times n$ matrices then $\det(AB) = (\det A)(\det B)$ and $\det(A^T) = \det A$.

Algorithm to compute $\det A$.

Input: an $n \times n$ matrix A .

1. Start by setting $\text{denom} = 1$.
2. Row reduce A to an echelon form E , while doing the following:
 - (a) When you switch two rows, multiply denom by -1 .
 - (b) When you rescale a row by a nonzero factor λ , multiply denom by λ .
 - (c) When you add a multiple of a row to another row, don't do anything to denom .

Then $\det A = \frac{\det E}{\text{denom}} = \frac{\text{the product of the diagonal entries of } E}{\text{denom}}$.

Here is one more way to compute $\det A$, which can be useful if there are many zero entries:

Theorem. Consider the matrix $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} & \cdots & a_{1n} \\ a_{21} & a_{22} & a_{23} & \cdots & a_{2n} \\ a_{31} & a_{32} & a_{33} & \cdots & a_{3n} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & a_{n3} & \cdots & a_{nn} \end{bmatrix}$.

Define $A^{(i,j)}$ as the submatrix formed by deleting row i and column j . Then

$$\det A = a_{11} \det A^{(1,1)} - a_{12} \det A^{(1,2)} + a_{13} \det A^{(1,3)} - \dots - (-1)^n a_{1n} \det A^{(1,n)}$$

Each $A^{(1,j)}$ is a square matrix smaller than A , so $\det A^{(1,j)}$ can be computed by the same formula.

Example. $\det \begin{bmatrix} a & b & c \\ d & e & f \\ g & h & i \end{bmatrix} = a(ei - fh) - b(di - fg) + c(dh - eg).$

This recursive formula for $\det A$ is most useful if A has many entries which are zero.

Example. If $A = \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 3 & 4 & 5 \\ 1 & 6 & 0 & 0 \\ 0 & 1 & 1 & 1 \end{bmatrix}$ then $\det A = \det \begin{bmatrix} 3 & 4 & 5 \\ 6 & 0 & 0 \\ 1 & 1 & 1 \end{bmatrix} - 0 + 2 \det \begin{bmatrix} 0 & 3 & 5 \\ 1 & 6 & 0 \\ 0 & 1 & 1 \end{bmatrix} - 0$ and

$$\det \begin{bmatrix} 3 & 4 & 5 \\ 6 & 0 & 0 \\ 1 & 1 & 1 \end{bmatrix} = \det \begin{bmatrix} 3 & 6 & 1 \\ 4 & 0 & 1 \\ 5 & 0 & 1 \end{bmatrix} = -\det \begin{bmatrix} 6 & 3 & 1 \\ 0 & 4 & 1 \\ 0 & 5 & 1 \end{bmatrix} = -\det \begin{bmatrix} 6 & 0 & 0 \\ 3 & 4 & 5 \\ 1 & 1 & 1 \end{bmatrix} = -6 \det \begin{bmatrix} 4 & 5 \\ 1 & 1 \end{bmatrix} = 6$$

since we taking transposes doesn't change the determinant, and switching columns reverses the sign of the determinant. Similarly, we have

$$\det \begin{bmatrix} 0 & 3 & 5 \\ 1 & 6 & 0 \\ 0 & 1 & 1 \end{bmatrix} = \det \begin{bmatrix} 0 & 1 & 0 \\ 3 & 6 & 1 \\ 5 & 0 & 1 \end{bmatrix} = -\det \begin{bmatrix} 3 & 1 \\ 5 & 1 \end{bmatrix} = -(3 - 5) = 2.$$

Therefore $\det A = 6 + 2 \cdot 2 = 10$.

2 Interpreting the determinant geometrically

The last thing we'll mention about determinants is something we explored in a demonstration last week:

Proposition. If A is an $n \times n$ matrix then $|\det A|$ is the volume of the n -dimensional parallelogram

$$P(A) = \{Av : v \in \mathbb{R}^n \text{ with } 0 \leq v_i \leq 1 \text{ for all } i = 1, 2, \dots, n\}.$$

Proof idea for $n = 2$ case. Assume $n = 2$ and $A = [u \ v]$ for some vectors $u, v \in \mathbb{R}^2$.

Make things simple by putting u and v both in the first quadrant. Draw a picture of the parallelogram $P(A)$ inside the rectangle R whose diagonal is $u + v$ and whose sides are on the x - and y -axes. Then compute the area of $P(A)$ by subtracting the areas of an appropriate number of rectangular and triangular regions from R . One finds that this area is $ad - bc$ if $u = \begin{bmatrix} a \\ c \end{bmatrix}$ and $v = \begin{bmatrix} b \\ d \end{bmatrix}$. \square

Corollary. Suppose $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is a linear transformation with standard matrix A . If S is any region in \mathbb{R}^n with finite volume then the volume of $T(S)$ is the volume of S times $|\det A|$.

Proof idea. If T is not invertible that $T(S)$ has zero volume (why?) while $\det A = 0$, so the result holds.

Assume T is invertible. Given a vector $u \in \mathbb{R}^n$ and a scalar $c > 0$, let $u + cS = \{u + cv : v \in S\}$.

Define $Q = \{v \in \mathbb{R}^n : 0 \leq v_i \leq 1 \text{ for all } i\}$. Then $P(A) = T(Q)$, so $T(u + cQ) = Au + cP(A)$.

The volume of $u + cQ$ is c^n and the volume of $T(u + cQ)$ is $c^n |\det A|$.

It follows that if $R \subseteq S$ is any disjoint union of translated rescaled cubes of the form $u + cQ$, then

$$\text{vol}(R)|\det A| = \text{vol}(T(R)) \leq \text{vol}(T(S)).$$

But $\text{vol}(S)$ is the limit superior of all such estimated volumes $\text{vol}(R)$, so $\text{vol}(S)|\det A| \leq \text{vol}(T(S))$.

The same argument with S replaced by $T(S)$ and T replaced by T^{-1} shows $\text{vol}(T(S))|\det A^{-1}| \leq \text{vol}(S)$.

Since $|\det A^{-1}| = 1/|\det A|$, it follows that $\text{vol}(S)|\det A| \geq \text{vol}(T(S))$ so $\text{vol}(S)|\det A| = \text{vol}(T(S))$. \square

3 Vector spaces

This course focuses on \mathbb{R}^n and its subspaces.

These objects are examples of (*real*) *vector spaces*.

There is also a notion of a *complex vector space* where our scalars can be complex numbers from \mathbb{C} rather than just \mathbb{R} . Essentially all of the theory is the same, so for now we stick to real vector spaces which are more closely aligned with applications.

The general definition of a vector space is given as follows:

Definition. A *vector space* is a nonempty set V with two operations called *vector addition* and *scalar multiplication* satisfying several conditions. We refer to the elements of V as *vectors*.

The vector addition operation for V must be a rule that takes two input vectors $u, v \in V$ and produces an output vector $u + v \in V$ such that

- (a) $u + v = v + u$.
- (b) $(u + v) + w = u + (v + w)$.
- (c) There exists a unique *zero vector* $0 \in V$ with the property that $0 + v = v$ for all $v \in V$.

The scalar multiplication operation for V must be a rule that takes a scalar input $c \in \mathbb{R}$ and an input vector $v \in V$ and produces an output vector $cv \in V$ such that

- (a) If $c = -1$ then $v + (-1)v = 0$.
- (b) $c(u + v) = cu + cv$.
- (c) $(c + d)v = cv + dv$ for $c, d \in \mathbb{R}$.
- (d) $c(dv) = (cd)v$ for $c, d \in \mathbb{R}$.
- (e) If $c = 1$ then $1v = v$.

Notation: If V is a vector space and $v \in V$ then we define $-v = (-1)v$ and $u - v = u + (-v)$.

Example. \mathbb{R}^n and any subspace of \mathbb{R}^n are vector spaces, with the usual operations of vector addition and scalar multiplication.

Example. The set $\mathbb{R}^{m \times n}$ of $m \times n$ matrices is a vector space. Its zero vector is the zero $m \times n$ matrix.

Example. The set of linear functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a vector space. Its zero vector is the function f with the formula $f(x) = 0$ for all $x \in \mathbb{R}^n$.

Example. The set of positive real numbers $\mathbb{R}^+ = \{x \in \mathbb{R} : x > 0\}$ is a vector space, but *not for the usual addition and multiplication operations*. Instead, define a new version of addition \oplus by $x \oplus y = xy$ for $x, y \in \mathbb{R}^+$ and new version of scalar multiplication \odot by $c \odot x = x^c$ for $c \in \mathbb{R}$ and $x \in \mathbb{R}^+$. The zero vector for this vector space $(\mathbb{R}^+, \oplus, \odot)$ is the number 1. We also have $\ominus x = 1/x$ and $x \ominus y = x/y$.

It is rarely necessary to check the axioms of a vector space in detail, and there is not much need to memorize the abstract definition. If we have a set with operations that look like vector addition and scalar multiplication for \mathbb{R}^n , then we usually have a vector space. Moreover, it's typical easy to identify every vector space we encounter as a special case of a few general constructions like the following:

Example. Let X be any set. Define $\text{Functions}(X, \mathbb{R})$ to be the set of functions $f : X \rightarrow \mathbb{R}$.

Given $f, g \in \text{Functions}(X, \mathbb{R})$ define $f + g$ to be the function with the formula

$$(f + g)(x) = f(x) + g(x) \quad \text{for } x \in X.$$

Given $c \in \mathbb{R}$ and $f \in \text{Functions}(X, \mathbb{R})$, define cf to be the function with the formula

$$(cf)(x) = cf(x) \quad \text{for } x \in X.$$

The set $\text{Functions}(X, \mathbb{R})$ is a vector space relative to these operations.

The corresponding zero vector in $\text{Functions}(X, \mathbb{R})$ is the function with the formula $f(x) = 0$ for all $x \in X$.

In a sense which can be made precise, the vector space \mathbb{R}^n “is the same as” $\text{Functions}(\{1, 2, 3, \dots, n\}, \mathbb{R})$.

More generally, if V is any vector space then the set of functions $\text{Functions}(X, V) = \{f : X \rightarrow V\}$ is a vector space for similar definitions of vector addition and scalar multiplication.

As an example of how one can use the axioms to prove properties of a general vector space, consider the following identities which are obvious for subspaces of \mathbb{R}^n .

Proposition. If V is a vector space then $0v = 0$ and $c0 = 0$ for all $c \in \mathbb{R}$ and $v \in V$.

Proof. We have $0v = (0 + 0)v = 0v + 0v$ so $0 = 0v - 0v = (0v + 0v) - 0v = 0v + (0v - 0v) = 0v + 0 = 0v$.

Similarly, $c0 = c(0 + 0) = c0 + c0$ so $0 = c0 - c0 = (c0 + c0) - c0 = c0 + (c0 - c0) = c0 + 0 = c0$. \square

We will not focus very much in this course on the art of coming up with these sorts of algebraic derivations. Mostly, we can just rely on our intuition from subspaces of \mathbb{R}^n when working with more general spaces.

4 Subspaces, bases, and dimension

Notions of subspaces, bases, and dimension for vector spaces are essentially the same as for \mathbb{R}^n .

Definition. A *subspace* of a vector space V is a subset H containing the zero vector of V , such that if $u, v \in H$ and $c \in \mathbb{R}$ then $u + v \in H$ and $cv \in H$.

If $H \subset V$ is a subspace then H is itself a vector space with the same operations of scalar multiplication and vector addition.

Example. V is a subspace of itself and $\{0\} \subset V$ is a subspace.

Example. \mathbb{R}^2 is technically not a subspace of \mathbb{R}^3 since \mathbb{R}^2 is not a subset of \mathbb{R}^3 .

If you want a subspace of \mathbb{R}^3 that “looks like” \mathbb{R}^2 , three candidates are

$$\left\{ \begin{bmatrix} x \\ y \\ 0 \end{bmatrix} : x, y \in \mathbb{R} \right\}, \quad \left\{ \begin{bmatrix} x \\ 0 \\ y \end{bmatrix} : x, y \in \mathbb{R} \right\}, \quad \text{and} \quad \left\{ \begin{bmatrix} 0 \\ x \\ y \end{bmatrix} : x, y \in \mathbb{R} \right\}.$$

Is there anything intrinsic that makes one of these subspaces more natural than the rest?

Example. Let X be any set. Let $Y \subset X$ be a subset. Define H as the subset of $\text{Functions}(X, \mathbb{R})$ consists of the functions $f : X \rightarrow \mathbb{R}$ with $f(y) = 0$ for all $y \in Y$. Then H is a subspace.

Example. The set of all functions $\text{Functions}(\mathbb{R}^n, \mathbb{R}^m)$ is a vector space since \mathbb{R}^m is a vector space. The subset of linear functions $f : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a subspace of this vector space.

Let V be a vector space.

Definition. A *linear combination* of a finite list of vectors $v_1, v_2, \dots, v_k \in V$ is a vector of the form

$$c_1v_1 + c_2v_2 + \dots + c_kv_k$$

for some scalars $c_1, c_2, \dots, c_k \in \mathbb{R}$. A linear combination of an infinite set of vectors is a linear combination of some finite subset. A linear combination by definition **only involves finitely many vectors**.

Definition. The *span* of a set of vectors is the set of all linear combinations that can be formed from the vectors. It is important to note that each such linear combination can **only involve finitely many vectors**. The span of a set of vectors in V is a subspace of V .

Example. Let $V = \text{Functions}(\mathbb{R}, \mathbb{R})$. The span of the infinite set of functions $1, x, x^2, x^3, \dots \in V$ is the subspace of *polynomial functions*. Each polynomial function is a linear combination of a finite number of monomials $c_nx^n + c_{n-1}x^{n-1} + \dots + c_1x + c_0$. An infinite sum like $1 + x + x^2 + \dots$ is **not** a polynomial.

Definition. A finite list of vectors $v_1, v_2, \dots, v_k \in V$ is *linearly independent* if it is impossible to express $0 = c_1v_1 + c_2v_2 + \dots + c_kv_k$ for some $c_1, c_2, \dots, c_k \in \mathbb{R}$ except when $c_1 = c_2 = \dots = c_k = 0$. An infinite list of vectors is linearly independent if every finite subset is linearly independent.

Definition. A *basis* of a vector space V is a subset of linearly independent vectors whose span is V . Saying that b_1, b_2, b_3, \dots is a basis for V is the same thing as saying that each $v \in V$ can be expressed as a uniquely linear combination of basis elements.

Theorem. Let V be a vector space.

1. V has at least one basis.
2. Every basis of V has the same size.
3. If A is a subset of linearly independent vectors in V then V has a basis B with $A \subset B$.
4. If C is a subset of vectors in V whose span is V then V has a basis B with $B \subset C$.

When V has a basis that is finite in size, the proof of the previous theorem is the same as for the case when V is a subspace \mathbb{R}^n (which was shown in earlier lectures). When V has no finite basis, the properties in the theorem still hold, but their proofs require some ideas beyond the scope of this course.

Definition. As for subspaces of \mathbb{R}^n , we define the *dimension* of a vector space V to be the common number of elements in any of its bases. Denote the dimension of V by $\dim V$.

Corollary. If $H \subset V$ is a subspace then $\dim H \leq \dim V$.

Moreover, if $H \subset V$ is a subspace with $\dim H = \dim V$ then $H = V$.

Proof. This follows from the last two parts of the previous theorem. □

Example. If X is a finite set then $\dim \text{Functions}(X, \mathbb{R}) = |X|$ where $|X|$ is the size of X . A basis is given

by the functions $\delta_y : X \rightarrow \mathbb{R}$ for $y \in X$, defined by the formulas

$$\delta_y(x) = \begin{cases} 1 & \text{if } x = y \\ 0 & \text{if } x \neq y \end{cases} \quad \text{for } x \in X.$$

Suppose U and V are vector spaces.

Definition. A function $f : U \rightarrow V$ is *linear* if

$$f(u + v) = f(u) + f(v) \quad \text{and} \quad f(cv) = cf(v) \quad \text{for all } c \in \mathbb{R} \text{ and } u, v \in U.$$

Define $\text{range}(f) = \{f(x) : x \in U\}$ and $\text{kernel}(f) = \{x \in U : f(x) = 0\}$.

Proposition. If $f : U \rightarrow V$ is linear then $\text{range}(f)$ and $\text{kernel}(f)$ are subspaces.

These subspaces are generalizations of the column space and null space of a matrix.

Proposition. If U, V, W are vector spaces and $f : V \rightarrow W$ and $g : U \rightarrow V$ are linear functions then $f \circ g : U \rightarrow V \rightarrow W$ is linear, where $f \circ g(x) = f(g(x))$.

Check this yourself!

5 Vocabulary

Keywords from today's lecture:

1. Vector spaces.

A vector space is a nonempty set V with two operations called *vector addition* and *scalar multiplication* that formally resemble the operations of vector addition and scalar multiplication for elements of \mathbb{R}^n . The precise definition involves a long list of axioms governing these operations, but in practice it's rarely necessary to remember the axioms.

Example: Any subspace of \mathbb{R}^n .

Example: Given a set X , the set $\text{Functions}(X, \mathbb{R})$ of functions $f : X \rightarrow \mathbb{R}$, provided we define $f + g$ as the function with the formula

$$(f + g)(x) = f(x) + g(x) \quad \text{for } x \in X$$

and define cf as the function with the formula

$$(cf)(x) = cf(x) \quad \text{for } x \in X$$

whenever $f, g : X \rightarrow \mathbb{R}$ and $c \in \mathbb{R}$.

2. Subspace of a vector space.

A nonempty subset closed under linear combinations.

3. Linearly combination and span of elements in a vector space.

A linear combination of a finite set of vectors $v_1, v_2, \dots, v_p \in V$ is a vector of the form

$$c_1v_1 + c_2v_2 + \dots + c_pv_p$$

where $c_1, c_2, \dots, c_p \in \mathbb{R}$. A linear combination of an infinite set of vectors is a linear combination of some finite subset. The set of all linear combinations of a set of vectors is the span of the vectors.

4. Linearly independent elements in a vector space.

A list of elements in a vector space is **linearly dependent** if one vector can be expressed as a linear combination of a finite subset of the other vectors. If this is impossible, then the vectors are linearly independent.

Example: $\cos(x)$ and $\sin(x)$ are linearly independently in $\text{Functions}(\mathbb{R}, \mathbb{R})$.

Example: the infinite list of functions $1, x, x^2, x^3, x^4, \dots$ are linearly independent in $\text{Functions}(\mathbb{R}, \mathbb{R})$.

5. Basis and dimension of a vector space.

A set of linearly independent elements whose span is the entire vector space.

Every basis in a vector space has the same number of elements. This number is defined to be the **dimension** of the vector space.

6. Linear functions.

If U and V are vector spaces, then a function $f : U \rightarrow V$ is linear when

$$f(u + v) = f(u) + f(v) \quad \text{and} \quad f(cv) = cf(v)$$

for all $u, v \in U$ and $c \in \mathbb{R}$.